REPORT E	Form Approved MB No. 3704-0188							
Number registration burgers and this is also constituted by the second of the constitute of the consti	consisting energy of this leasesthing as some das a vousing distance of the second data would be seen as the second definition of the second design of the s	-						
Tillings of the Miles of the Control								
1. AGENCY USE ONLY (Leave on	January 12, 199		D DATES COVERED  # 29					
Structural, Magn	etic and Electrical ry CePdIn <sub>2</sub>	Properties	S. WHOMG HUMBERS N00014-93-1-0904					
5. AUTHOR(S)	Dr. John Pazik							
Y. Ijiri, F.J. D	iSalvo and H. Yamar	ie	R&T: 3134037ess08					
7. PERFORMING ORGANIZATION	NAME(S) AND ADDRESS(ES)		8. FERFORMING ORGANIZATION REPORT NUMBER					
Department of Ch Cornell Universi Ithaca, NY 1485	ty							
3. SPONSORING/MONITORING A	SENCY NAME(S) AND ADDRESS(ES		10. SPONSORING / MONITORING GENCY PEPORT NUMBER					
Office of Naval Chemistry Progra 800 N. Quincy St Alexandria, VA	m •	1996	0205044 —					
11. SUPPLEMENTARY MOTES			2000 OTI -					
	statement s been approved fo: ; its distribution		12b. DISTRIBUTION CODE					
Cmcm (Z=4) with a crystal X-ray dat independent refle BRe3 structure ty Magnetic measurem phase transition	ound to crystallize $=4.6210(9)$ Å, b=10.6 a were refined to Ections and 16 variate and is the first ents indicate that with a $T_{\rm C}$ near 10K.	594(2)Å, and c R=0.0459 (wR2= ables. The st known Ce-con CePdIn <sub>2</sub> under The electri	0.0666), with 255 ructure adopts the	>				
14. SUBJECT TERMS			15. NUMBER OF PAGES					
·			16. PRICE CODE					
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFIC OF ABSTRACT	CATION 20. LIMITATION OF ABSTRA	\CT				
Unclassified	Unclassified	Unclassifie	ed					

#### OFFICE OF NAVAL RESEARCH

Grant or Contract N00014-93-1-0904

R&T Code 3134037ess08 Scientific Officer: Dr. John Pazik

Technical Report No. 29

"Structural, Magnetic and Electrical Properties of the New Ternary CePdIn2"

bу

Y. Ijiri, F.J. DiSalvo and H. Yamane

Submitted to

J. Solid State Chemistry

Cornell University
Department of Chemistry
Ithaca, NY 14853

January 12, 1996

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

Nevised Version 11/6/95

# Structural, Magnetic and Electrical Properties of the New Ternary CePdIn<sub>2</sub>

Y. Ijiri and F.J. DiSalvo\* Department of Chemistry, Cornell University, Ithaca, New York 14853

H. Yamane Institute for Materials Research, Tohoku University, Sendai (Japan)

\*Author to whom correspondence should be addressed.

#### **ABSTRACT**

CePdIn<sub>2</sub> is found to crystallize in the orthorhombic space group *Cmcm* (Z=4) with a = 4.6210(9) Å, b = 10.694(2) Å, and c = 7.455(2) Å. Single crystal X-ray data were refined to R = 0.0459 (wR<sub>2</sub> = 0.0666), with 255 independent reflections and 16 variables. The structure adopts the BRe<sub>3</sub> structure type and is the first known Ce-containing member. Magnetic measurements indicate that CePdIn<sub>2</sub> undergoes a ferromagnetic phase transition with a T<sub>c</sub> near 10 K. The electrical resistivity of CePdIn<sub>2</sub> shows metallic behavior with a small anomaly at the magnetic phase transition.

#### INTRODUCTION

Over the past 20 years, research in ternary cerium intermetallics has grown substantially as interesting strongly correlated electron systems have been observed (1,2,3). While much progress has been made in understanding the origin of these effects, this knowledge is insufficient to predict new compounds which exhibit these phenomena.

However, as discussed by Fisk et al. (4), the chemical composition of the interesting Ce compounds includes mostly elements in the late transition metal and early p block columns. In particular, in the Ce-Pd-In ternary phase diagram, there are a number of interesting correlated electron materials such as CePd<sub>3</sub> (5), CeIn<sub>3</sub> (6), and CePdIn (7).

Motivated by the unusual materials found in this system as well as more recently discovered compounds (8), we have been searching for other new ternaries in this phase space (9). Here, we report the structure and properties of a new compound, CePdIn<sub>2</sub>.

#### **EXPERIMENTAL**

Sample Preparation

CePdIn<sub>2</sub> was synthesized by arc-melting stoichiometric amounts of the elements on a water-cooled, Ta coated copper hearth under a flow of Tigettered Ar (Centorr Furnaces, Model 2B-20). The arc-melted beads were flipped over and remelted repeatedly to increase homogeneity. Mass losses after arc-melting were less than 0.5%.

The elements used were all at least of purity 99.9%. The cerium was further purified by vacuum melting into a water-cooled copper cup, leaving behind much of the outer oxide coat often present on Ce pieces.

The arc-melted beads were then placed in tantalum tubing and sealed in quartz under vacuum. The beads were then heated at 825 °C for one week. Structure determination

Preliminary structure characterization was done with electron and x-ray diffraction of finely ground powder. Electron diffraction patterns were obtained using powder suspended on holey carbon coated grids in a double tilt holder in a JEOL 1200 EX transmission electron microscope. The powder was also studied by X-ray diffraction using a Scintag XDS 2000 diffractometer with Cu  $K_{\alpha}$  radiation. The diffraction data indicated an orthorhombic cell of approximate dimensions 4.621 x 10.692 x 7.450 Å and C-centering.

Shards of lightly cracked beads were mounted and screened for crystal quality using precession camera photos. Single crystal data were collected on a Siemens P4 four-circle diffractometer in the  $\theta$ -2 $\theta$  mode from -1  $\leq$  h  $\leq$  6, -1  $\leq$  k  $\leq$  13, and -1  $\leq$  1  $\leq$  9. The lattice constants for the cell were a = 4.6210(9) Å, b = 10.694(2) Å, and c = 7.455(2) Å, in good agreement with the powder data. Three check reflections were measured every 50 reflections, and they showed no evidence of crystal decomposition or movement on the fiber. An empirical psi-scan absorption correction was applied to the data, yielding an  $R_{int}$  = 0.0305. The semi-invariants representation program SIR92 (10) was used to generate a model for the structure which was refined with XL from the SHELXTL

version 5 $\gamma$  program (11). The final structure R values were wR<sub>2</sub> = 0.0666 and R<sub>1</sub> = 0.0459 for all data and wR<sub>2</sub> = 0.0614 and R<sub>1</sub> = 0.0310 for I > 2 $\sigma$ (I). The MISSYM algorithm (12) detected no further symmetry elements than those expected for Cmcm.

Crystal structure data, values of the atomic parameters, and anisotropic displacement parameters are listed in Tables 1, 2, and 3. The atomic positions are presented both in accordance with previous BRe<sub>3</sub> structure literature and also in the standard format as determined by the program STRUCTURE TIDY (13). This program standardizes inorganic structures to help identify similar structures, obscured by differences in the reporting of crystal data.

Additional structure information, including structure amplitudes, is available as supplementary material.<sup>1</sup>

Physical property measurements

The magnetic susceptibility of CePdIn<sub>2</sub> was measured using a Faraday balance on coarsely ground, loose powder. The field dependence of the susceptibility at room temperature from 2.1 kG to 14.6 kG showed variations of less than 1%, indicating the absence of ferromagnetic impurities. Temperature dependent data from 4 to 325 K were collected in a field of 10 kG. Platinum was used as a calibration standard. The measured room temperature susceptibility of Pt  $\chi$ (293 K) was 1.030 x 10<sup>-6</sup> emu/g, in good agreement with a literature value of 1.035 x 10<sup>-6</sup> emu/g (14).

Electrical resistivity was measured using standard four probe AC techniques. The contacts were ohmic by linearity of the I-V curve. The

resistivity sample was bar-shaped, cut with a string saw from a bead after heat treatment.

#### **RESULTS AND DISCUSSION**

Structural Aspects

The structure of CePdIn<sub>2</sub> is isotypic with the rare earth intermetallics (Y,Gd, Tb, Dy)NiIn<sub>2</sub> (15), (Y,Tb-Lu)NiGa<sub>2</sub> (16), (Y, Tb-Lu)PdGa<sub>2</sub> (17), and (Y, Tb-Lu)NiAl<sub>2</sub> (18). Thus, the structure type was previously known only for compounds containing the smaller rare earth atoms.

As shown in Figure 1, the structure can be described as a staggered net of trigonal prisms. Each prism is composed of four In atoms and two Ce atoms centered by a Pd atom.

Selected bond distances are listed in Table 4. The bond distances indicate that the chemical environment of the atoms is more complicated than emphasized in Figure 1. Figure 2 shows the coordination polyhedra around each of the atoms as determined by the principle of maximal gap in near neighbor distances (19). All resulting distances are short enough to be considered as bonding to the central atom.

As shown in Figure 2a, the 13-coordinate environment around the Ce can be described as a distorted pentagonal prism of eight In atoms and two Pd atoms with two In atoms and one Pd atom capping three square faces of the prism. The Pd environment of nine neighbors, shown in Figure 2b, consists of a trigonal prism of four In atoms and two Ce atoms, tricapped by two In

atoms and one Ce atom. The coordination geometry of In to its 12 neighbors is irregular and is depicted in Figure 2c. It consists roughly of Ce atoms in square pyramidal coordination and In atoms in tetrahedral coordination about the central In. Three Pd atoms complete the coordination polyhedron.

The complicated polyhedra suggest that size constraints strongly limit the formation of the BRe<sub>3</sub> structure type. An unsuccessful attempt at making CeNiIn<sub>2</sub> underscores this probable steric limitation. The larger Pd and In atoms are perhaps necessary to stabilize the structure for the larger rare earth Ce atom. To date, the presented structure is the only known Ce or Pd-In example for this structure type.

### Magnetic properties

Figure 3 shows the inverse magnetic susceptibility for CePdIn<sub>2</sub> upon heating the powder sample from 4.2 K. The data were fit to the Curie-Weiss law:  $\chi = \chi_0 + \frac{C}{T-\theta}$ .

The fit parameters were selected by choosing a temperature range of fit, fixing  $\theta$ , and then determining C and  $\chi_0$  using standard analysis techniques (20).  $\theta$  was varied, and for each choice, a fractional variance  $\sigma$  was calculated. The final parameters were chosen to minimize  $\sigma$  over the largest temperature range.

These parameters were found to be  $\chi_0 = 2.6 \pm 0.4 \times 10^{-7}$  emu/g, C = 1.60  $\pm$  0.02  $\times$  10<sup>-3</sup> emu-K/g, and a ferromagnetic  $\theta = 16.5 \pm 1.5$  K for the temperature

range of 75 to 325 K with a  $\sigma$  = 1.5 x 10<sup>-3</sup>. At temperatures below 75 K, some deviations from the Curie-Weiss law occur, most likely due to crystal field effects. The Curie constant corresponds to a moment of 2.47  $\pm$  0.02  $\mu_B$  which is very close to the free ion value of 2.54  $\mu_B$  for Ce<sup>+3</sup>. Hence, it appears that in CePdIn<sub>2</sub>, the Ce is trivalent, with ferromagnetic exchange. We can estimate a transition temperature of  $\approx$  10 K from the inflection point of the  $\chi$  vs temperature curve.

As shown in Figure 4, the field dependence of the low temperature susceptibility is consistent with a ferromagnetically ordered material. Fitting the  $\chi$  vs 1/H curve to a straight line, we obtain a magnetization at 4.2 K of 1.6  $\mu_B$  vs the expected gJ = 2.14  $\mu_B$  for completely oriented Ce<sup>+3</sup> ions in a J=5/2 state. The deviation from the expected value may indicate that there is some canting of the spins in the ordered state or more likely that crystal field splitting reduces the ground state moment.

The existence of ferromagnetism in this system agrees well with the empirical observations of Sereni and Kappler (21). They found that ferromagnetic behavior is often present in systems with Ce-Ce nearest neighbor distances between 3.7 and 4.1 Å. In CePdIn<sub>2</sub> this distance is 4.063 Å. Electrical properties

Resistivity data from room temperature to 4.2 K are shown in Figure 5. The magnitude of the resistivity and its monotonic decrease with temperature down to  $\approx 20$  K indicate metallic conduction. At roughly 10 K,

there is a kink in the resistivity, most likely due to a coupling of the carriers to the ordering Ce moments.

Note that the resistivity decreases more rapidly below 10 K than just above, indicating that spin-disorder scattering may be playing a role at higher temperatures. Below the ferromagnetic transition, this extra scattering is expected to die out as the moments order. This behavior is similar to that observed in other rare earth intermetallic compounds such as NdAuAl (22) and GdNiGe (23).

#### CONCLUSIONS

We have synthesized a new ternary intermetallic compound,  $CePdIn_2$ , the first known Ce containing member of the  $BRe_3$  structure type. From the magnetic and electrical properties, we conclude that  $CePdIn_2$  is a ferromagnet with a  $T_c$  of about 10 K. It appears that the Ce moments do not interact strongly with the conduction electrons, as we see little evidence of Kondo-like magnetic or transport behavior.

# **ACKNOWLEDGMENTS**

This work was supported by the Office of Naval Research. We would like to thank E. Karr for her help with the electron diffraction and N.E. Brese for his help with the x-ray crystallography. Y. I. would also like to acknowledge financial support from a National Physical Science Consortium fellowship.

#### **REFERENCES**

- 1. Z. Fisk and J.R. Schrieffer, MRS Bulletin 28, 23 (1993).
- 2. N.B. Brandt and V.V. Moshchalkov, Adv. Phys. 33, 373 (1984).
- 3. J.M. Lawrence and D.L. Mills, Comments Cond. Mat. Phys. 15, 163 (1991).
- 4. Z. Fisk, J.D. Thompson, and H.R. Ott, J. Magn. Magn. Mater. 76&77, 637 (1988).
- 5. D. Jaccard, M.J. Besnus, and J.P. Kappler, J. Magn. Magn. Mater. 63&64, 572 (1987).
- 6. J.M. Fournier and E. Gratz in K.A. Gschneidner, Jr. L. Eyring, G. H. Lander and G.R. Choppin (eds.), Handbook on the Physics and Chemistry of Rare Earths, Elsevier, Netherlands, 17, 409 (1993).
- 7. K. Satoh, T. Fujita, Y. Maeno, Y. Uwatoko, and H. Fujii, *J. Phys. Soc. Japan* **59**, 692 (1990).
- 8. B. Xue, F. Hulliger, Ch. Baerlocher, and M. Estermann, J. Alloys Compd. 191 L9 (1993).
- 9. R.A. Gordon, Y. Ijiri, C.M. Spencer, and F.J. DiSalvo, J. Alloys Compd. 224, 101 (1995).
- 10. A. Altomere, M.C. Burla, M. Camilli, G. Cascarano, C. Giacovazzo, A. Guagliardi, G. Polidori, *J. Appl. Crystallogr.*, in preparation, (1994).
- 11. G.M. Sheldrick, SHELXTL version 5, Siemens Analytical X-ray Instruments, Inc., Madison, WI.

- 12. Y. LePage, J. Appl. Crystallogr. 20, 264 (1987).
- 13. L.M. Gelato and E. Parthé, J. Appl. Crystallogr. 20, 139 (1987).
- 14. D.R. Lide, ed., Handbook of Chemistry and Physics, CRC Press, Boca Raton, 74, 9-53 (1993).
- 15. V.I. Zaremba, O.Ya. Zakharko, Ya.M. Kalychak, and O.I. Bodak, Dop. Akad.
- Nauk Ukrain. RSR, Seriya B 12, 44 (1987).
- 16. V.A. Romaka, Ju.N. Grin', Ja.L. Jarmoljuk, R.V. Skolozdra, and A.A. Jartys', Ukr. Fiz. Zh. 28, 227 (1983).
- 17. Ju.N. Grin', Dop. Akad. Nauk Ukrain. RSR, Seriya B 9, 34 (1984).
- 18. R.M. Rykhal', O.S. Zarechnyuk, and Ya. P. Yarmolyuk, Sov. Phys. Cryst. 17, 453 (1972).
- 19. G.O. Brunner and D. Schwarzenbach, Z. Kristallogr. 133, 127 (1971).
- 20. P.R. Bevington, Data Reduction and Error Analysis for the Physical Sciences, McGraw Hill, New York, (1969).
- 21. J.G. Sereni and J.P. Kappler, *Physica B* **171**, 166 (1991).
- 22. L. Menon and S.K. Malik, Phys. Rev. B 51, 5858 (1995).
- 23. C.S. Garde, J. Ray, and G. Chandra, J. Alloys Compd. 201, L5 (1993).

TABLE 1  $\label{eq:crystal} \mbox{Crystal Data and Structure Refinement for CePdIn}_2$ 

Tourisies I formula	CoDdin
Empirical formula	CePdIn <sub>2</sub>
Formula weight	476.16 Siemens P4 four-circle
Diffractometer type	
Radiation	Graphite monochromated $MoK_{\alpha}$ ,
	λ=0.71073 Å
Temperature	293(2) K
Scan type	θ-2θ
Crystal system	Orthorhombic
Space group	Cmcm (No. 63)
Unit cell dimensions	a = 4.6210(9) Å
	b = 10.694(2)  Å
	c = 7.455(2)  Å
Volume	$368.40(13) \text{ Å}^3$
Z	4
Density (calculated)	8.585 g/cm <sup>3</sup>
Absorption coefficient	29.012 mm <sup>-1</sup>
F(000)	808
Crystal size	$0.19 \times 0.05 \times 0.04 \text{ mm}$
$\theta$ range for data collection	3.81° to 27.48°
Index ranges	$-1 \le h \le 6$ , $-1 \le k \le 13$ , $-1 \le l \le 9$
Reflections collected	391
Independent reflections	$255 (R_{int} = 0.0305)$
Absorption correction	Semi-empirical from psi-scans
Max. and min. transmission	0.984 and 0.509
Refinement method	Full-matrix least-squares on F <sup>2</sup>
Data/restraints/parameters	255/0/16
Goodness-of-fit on F <sup>2</sup>	1.089
Final R indices (I>2 $\sigma$ (I))	$R_1 = 0.0310$ , $wR_2 = 0.0614$
R indices (all data)	$R_1 = 0.0459$ , $wR_2 = 0.0666$
Largest diff. peak and hole	2.547 and -1.546 eÅ <sup>-3</sup>

TABLE 2

Atomic Coordinates' and Equivalent Isotropic
Displacement Parameters (Å<sup>2</sup> x 10<sup>3</sup>) for CePdIn<sub>2</sub>

	x	y	Z	U(eq)
In in 8(f)	0.0 [0.0]	0.3612(1) [0.1388]	0.0506(1) [0.0506]	11(1)
Ce in $4(c)$	[0.0] 0.0	0.0756(1) [0.4244]	0.25 [0.25]	10(1)
Pd in 4(c)	[0.0]	0.7931(2) [0.7069]	0.25 [0.25]	11(1)

<sup>\*</sup>Coordinates in brackets are in STRUCTURE TIDY format.

TABLE 3  $\label{eq:Anisotropic Displacement Parameters ($\mathring{A}^2$ x 10³) for CePdIn}_2$ 

	$U_{11}$	$U_{22}$	$U_{33}$	$U_{23}$	$U_{13} = U_{12}$
Pd	9(1)	12(1)	10(1)	0	0
Ce	10(1)	11(1)	11(1)	0	0
In	11(1)	13(1)	8(1)	1(1)	0

TABLE 4

Interatomic Bond Distances (Å)

for CePdIn<sub>2</sub>


Ce	1Pd	3.021(3)
	<b>2P</b> d	3.278(2)
	4In	3.2891(9)
	2In	3.396(2)
	4In	<b>3.</b> 5786(13)
Pd	1Ce	3.021(3)
	<b>2</b> Ce	3.278(2)
	2In	2.783(2)
	4In	<b>2.8421(9)</b>
In	<b>2</b> Ce	3.2891(9)
	1Ce	3.396(2)
	<b>2</b> Ce	3.5786(13)
	1Pd	2.783(2)
	2Pd	2.8421(9)
	1In	2.973(2)
	1In	3.064(2)


#### LIST OF FIGURE CAPTIONS


- FIG. 1. Structure of CePdIn<sub>2</sub> illustrating trigonal prismatic coordination around the Pd atoms.
- FIG. 2. Local coordination environments around a) Ce b) Pd and c) In atoms. The dark lines indicate the pentagonal and trigonal prismatic units in the Ce and Pd coordination polyhedra.
- FIG. 3. Inverse magnetic susceptibility as a function of temperature. Inset shows susceptibility vs temperature for T < 50 K.
- FIG. 4. Magnetic susceptibility vs inverse field at 4.2 K.
- FIG. 5. Electrical resistivity as a function of temperature. Inset shows resistivity for T < 25 K.


#### LIST OF FOOTNOTES


<sup>1</sup> See NAPS document No. \_\_ for \_\_ pages of supplementary material. Order from ASIS/NAPS. Microfiche Publications, P.O. Box 3513, Grand Central Station, New York, NY 10163. Remit in advance \$4.00 for microfiche copy or for photocopy, \$7.75 up to 20 pages plus \$0.30 for each additional page. All orders must be prepaid. Institutions and Organizations may order by purchase order. However, there is a billing and handling charge for this service of \$15. Foreign orders add \$4.50 for postage and handling, for the first 20 pages, and \$1.00 for additional 10 pages of material, \$1.50 for postage of any microfiche orders.

















# Supplementary information: observed and calculated structure factors

							100- 1	1024 1	10-	b	le .	1 10F	o 10Fc 1	0=	p x	7	TOLO T	are To	78
	1 10Fo 10Fc 10s	h k	1 10Fo	10Fc 10s	h k	1	10Fo 1	LOPC .	108	-	-								
n K	1 1010 1010 100									_	_		4 381	60	1 3	6	319	90 4	15
			1 0	79 1.	1 11	2	769	764	28	4	•	4 40			3 3	6	153	65 13	53
2 0	0 6254 6015 40	5 5		• • • • • • • • • • • • • • • • • • • •	3 11		590	637	64	1	5	4 30		56		_			37
4 0	0 3912 3868 31	0 6	1 281				628	630	47	3	5	4 43	6 155	74	0 4	6			
	0 2435 2462 24	26	1 0	28 1	0 12					5	5	4 33	4 116	97	24	- 6	479	503	72
• •	U 2000	4 6	1 0	27 1	2 12			575	60					37	4 4	6	402	377 1	82
1 1	V		1 679	689 17	1 13	2	933	875	27	0	-			-	1 5	6	195	213 13	23
3 1	0 135 62 134	_		546 48	1 1		2292	2277	20	2	6	4 78	•	41		6		176 1	
5 1	0 171 99 98	37	1 539			-	1628		26	4	6	4 61	0 583	50	3 5				
7 2	0 858 834 16	57	1 346		3 1				20	i	7	4 240	5 2443	26	0 6	6			41
	•	0 8	1 400	407 44	5 1		1046		_	_	-			32	26	6	662	631	53
2 2		2 8	1 311		0 2	3	3103	3041	39	3	•			35	1 7	6	2079	2072	45
4 2	0 408 375 46		_		2 2	3	2481	2567	42	0	8	4 91	-		- :		1695		36
1 3	0 1873 1910 39	4 8	1 218				1785		29	2	8	4 88	5 846	39	3 7	_			
3 3	0 1329 1353 31	1 9	1 185		-				21	4	8	4 61	7 669	60	0 8	6			44
		39	1 284	116 182	1 3		1286			ī	_	4 28	6 65	74	2 8	- 6	685	646	60
5 3	Q 401	0 10	1 1646	1715 26	3 3		995		25		-			248	1 9	6	0	17.	1
0 4	0 2752 2798 45		1 1556		5 3	3	764	733	34		-				0 10	6	300	249 1	22
2 4	0 2315 2318 38	2 10			0 4	3	1789	1806	43	0	10	4 35		82	:				15
4 4	0 1568 1556 32	4 10	1 1261				1475		44	2	10	4 42	7 277	81	1 1	-	1353		
_	0 966 970 22	1 11	1 1527	1554 16	2 4					1		4 67	6 679	34	3 1		1111		22
	• • • • • • • • • • • • • • • • • • • •	3 11	1 1376	1320 30	4 4		1135		31			4 2		142	0 2	7	1750	1824	31
35		0 12	1 303		1 5	5 3	2492	2532	31	0				17	2 2		1682		38
5 5	0 540 444 42	-			3 :	. 3	1923	1967	35	1	1		7 1361				1262		33
0 6	0 158 146 158	2 12	1 320		_		1328		30	3	1	5 10	9 1062	20	4 2				
2 6	0 215 150 120	1 13	1 66						44	5	ī	5 7	6 717	27	1 3			845	24
_		0 0	2 866	852 19	0 (		2278			-			8 1677	40	3 3	7	680	725	56
4 6		2 0	2 715	692 18	2 (	63	1933	1927	45	0	_			39	0 4	. 7	1116	1127	26
1 7	0 4454 4472 31	_			4 (	6 3	1381	1401	30	2	-		8 1446		-		1059		40
37	0 3495 3509 32	4 0	2 533	,	i				116	4	2		1025	27		_			25
5 7	0 2437 2453 25	1 1	2 372			-		183	1	1	3	5 29	94 2981	38	1 !		1668		
	• • • • • • • • • • • • • • • • • • • •	3 1	2 2702	2 2604 29	_	7 3	-	-	_	3	3		32 2390	33	3 9	, 7	1426	1385	38
0 8	•	5 1	2 1720	5 1707 23	0 1	3	953	956	32	_	_			38	0 (	5 7	1417	1421	30
2 8	0 217 26 167	_			2 1	8 3	873	846	39	0	4		92 2741		-		1314		39
4 8	0 0 38 1	0 2			_	B 3	683	638	47	2	4		19 2443	42					
1 9	0 176 51 176	22	2 280		-		1738		26	4	4	5 18	09 1847	26	1 1			177	
	0 151 40 150	4 2	2 (	D 186 l						ĭ	5		32 1182	60	0 1	3 7	663	677	50
3 9	·	1 3	2 274	2738 29	3 9	9 3	1389		33				21 929	38	2	3 7	675	627	64
0 10	0 020			1 1966 37	0 1	0 3	503	469	55	3	5				_		1251	1219	36
2 10	0 438 470 63	3 3			2 1	0 3	345	438	107	٥	6	5 10	86 1104	30	_			322	
4 10	0 238 341 238	5 3				-	1182		57	2	6	5 9	36 <b>9</b> 82	42	_	0 1			
1 11	0 1742 1784 22	0 4	2 221	9 2216 46	1 1				38	4	6	5 7	40 726	48	2		3 229		200
	• • • • • • • • • • • • • • • • • • • •	2 4	2 183	8 1848 46	3 1		1100			_	7		42 726	26	1	1 1	8 1811	1802	12
3 11	• ••••	4 4		4 1246 30	0 1		1430		29	1		_	:	56	3	1 1	1513	1524	22
0 12	0 589 551 46	-			2 1	2 3	1257	1278	36	3	7							171	81
2 12	o 551 507 56	1 5	_		īī		1280		84	0	8		59 1036	32				153	ī
1 13	0 276 184 136	35					3625		36	2	8	5 9	55 937	42		2 3			_
	1 236 187 14	5 5		5 343 116	-					ī	9	5 11	03 1105	23	1		8 1374		20
		0 6	2 231	5 2295 46	2		6 3131			3	_		90 929	49	3	3	8 1152	1181	47
3 1	• • •	2 6		9 2005 45	4	0 4	4 2241	2202		_				27	ō	4	B 1099	1156	30
5 1	1 90 105 89			2 1465 30	1	1 4	4 1210	1166	25	-	10		71 1982		-		8 1064		45
0 2	1 232 251 42	4 6	-				4 952			2	10		29 1825	35	_				48
2 2	1 228 175 35	1 7	-		_	_				1	11	5 14	48 1428	24	_		8 417		
	1 127 70 127	3 7	2 47			_	- ::::			ō			52 2806	33	0		8 1291		31
		. 5 7	2 29	6 366 120		_	4 191	-		_			93 2513	41	2	6	8 1156	1180	48
1 3		_		6 2685 41	2	2	4 240			2					_	_	8 395	257	56
3 3	1 2397 2345 25						4 94	136	94	4	0		10 1915	24					64
5 3	1 1587 1566 17	2 8		,	_		4 0			1	1	6 1	858	12		_			
0 4	1 3137 3244 32	4 8	2 175	3 1808 28	_	_	-	28	_	3	_	6	112 718	24	0	2	9 0	35	. 1
	1 2566 2699 28	1 9	2	0 47 1	_	-	4 0	- :	_	_			37 175	65	2	2	9 355	34	124
. 2 4		3 9		7 39 197	5	3	4 191			0		-		151	ī	3	9 1290	1318	21
4 4				4 1693 38	ō	4	4 590	602	36	2					-	_	9 1342		30
1 5	1 174 71 50	0 10			_	_	4 572		49	4	2	6	139 117	139	0	4	7 1344	1333	
	4 901 94 78	2 10	2 156	0 1536 34	4	-	/-			_									

a. m3#1	2.783(2)	In-Pd#2	2.8421(9)
In-Pd#1 In-Pd#3	2.8421(9)	In-In#4	2.973(2)
In-Pa#5 In-In#5	3.064(2)	In-Ce#6	3.2891(9)
	3.2891(9)	In-Ce#8	3.396(2)
In-Ce#7 In-Ce#2	3.5786(13)	In-Ce#3	3.5786(13)
	3.021(3)	Ce-Pd#9	3.278(2)
Ce-Pd Ce-Pd#10	3.278(2)	Ce-In#11	3.2891(9)
Ce-In#12	3.2891(9)	Ce-In#13	3.2891(9)
Ce-In#12 Ce-In#14	3.2891(9)	Ce-In#15	3.396(2)
Ce-In#8	3.396(2)	Ce-In#2	3.5786(13)
Ce-In#16	3.5786(13)	Ce-In#17	3.5786(13)
Pd-In#18	2.783(2)	Pd-In#4	2.783(2)
Pd-In#17	2.8421(9)	Pd-In#2	2.8421(9)
Pd-In#3	2.8421(9)	Pd-In#16	2.8421(9)
Pd-Ce#19	3.278(2)	Pd-Ce#20	3.278(2)
Pa-Ce#19	30273(=,		
Pd#1-In-Pd#2	105.63(3)	Pd#1-In-Pd#3	105.63(3)
Pd#2-In-Pd#2	108.77(5)	Pd#1-In-In#4	143.65(4)
Pd#2-In-In#4	58.47(2)	Pd#3-In-In#4	58.47(2)
Pd#1-In-In#5	112.10(6)	Pd#2-In-In#5	112.15(5)
Pd#3-In-In#5	112.15(5)	In#4-In-In#5	104.26(4)
Pd#1-In-Ce#6	64.73(4)	Pd#2-In-Ce#6	168.55(4)
Pd#3-In-Ce#6	80.67(2)	In#4-In-Ce#6	132.95(2)
In#5-In-Ce#6	68.46(3)	Pd#1-In-Ce#7	64.73(4)
Pd#2-In-Ce#7	80.67(2)	Pd#3-In-Ce#7	168.55(4)
In#4-In-Ce#7	132.95(2)	In#5-In-Ce#7	<b>68.4</b> 6(3)
Ce#6-In-Ce#7	89.25(3)	Pd#1-In-Ce#8	79.60(4)
Pd#2-In-Ce#8	62.67(4)	Pd#3-In-Ce#8	62.67(4)
In#4-In-Ce#8	64.05(2)	In#5-In-Ce#8	168.30(6)
Ce#6-In-Ce#8	118.89(3)	Ce#7-In-Ce#8	118.89(3)
Pd#1-In-Ce#2	135.59(2)	Pd#2-In-Ce#2	54.69(5)
Pd#3-In-Ce#2	118.15(4)	In#4-In-Ce#2	65.46(2)
In#5-In-Ce#2	58.75(3)	Ce#6-In-Ce#2	127.21(3)
Ce#7-In-Ce#2	72.40(2)	Ce#8-In-Ce#2	113.23(3)
Pd#1-In-Ce#3	135.59(2)	Pd#2-In-Ce#3	118.15(4)
Pd#3-In-Ce#3	54.69(5)	In#4-In-Ce#3	65.46(2)
In#5-In-Ce#3	58.75(3)	Ce#6-In-Ce#3	72.40(2)
Ce#7-In-Ce#3	127.21(3)	Ce#8-In-Ce#3	113.23(3)
Ce#2-In-Ce#3	80.43(4)	Pd-Ce-Pd#9	135.19(3)
Pd-Ce-Pd#10	135.19(3)	Pd#9-Ce-Pd#10	89.62(6)
Pd-Ce-In#11	101.87(3)	Pd#9-Ce-In#11	50.14(2)
Pd#10-Ce-In#11	110.44(4)	Pd-Ce-In#12	101.87(3)
Pd#9-Ce-In#12	110.44(4)	Pd#10-Ce-In#12	50.14(2)
In#11-Ce-In#12	156.27(6)	Pd-Ce-In#13	101.87(3)
Pd#9-Ce-In#13	50.14(2)	Pd#10-Ce-In#13	110.44(4)
In#11-Ce-In#13	85.90(3)	In#12-Ce-In#13	89.25(3)
Pd-Ce-In#14	101.87(3)	Pd#9-Ce-In#14	110.44(4)
Pd#10-Ce-In#14	50.14(2)	In#11-Ce-In#14	89.25(3)
In#12-Ce-In#14	85.90(3)	In#13-Ce-In#14	156.27(6)
Pd-Ce-In#15	154.05(2)	Pd#9-Ce-In#15	50.36(3)
Pd#10-Ce-In#15	50.36(3)	In#11-Ce-In#15	61.11(3)
In#12-Ce-In#15	96.51(3)	In#13-Ce-In#15	96.51(3)
In#12-Ce-In#15	61.11(3)	Pd-Ce-In#8	154.05(2)
Pd#9-Ce-In#8	50.36(3)	Pd#10-Ce-In#8	50.36(3)
In#11-Ce-In#8	96.51(3)	In#12-Ce-In#8	61.11(3)
TESTT-CA-THEO			

- *** - ****	61 11/3)	In#14-Ce-In#8	96.51(3)
In#13-Ce-In#8	61.11(3) 51.91(4)	Pd-Ce-In#2	50.15(2)
In#15-Ce-In#8	• •	Pd#10-Ce-In#2	89.98(3)
Pd#9-Ce-In#2	155.46(2)	In#12-Ce-In#2	52.79(3)
In#11-Ce-In#2	150.26(4)		87.78(2)
In#13-Ce-In#2	107.60(2)	In#14-Ce-In#2	113.23(3)
In#15-Ce-In#2	139.28(2)	In#8-Ce-In#2	• •
Pd-Ce-In#16	50.15(2)	Pd#9-Ce-In#16	155.46(2)
Pd#10-Ce-In#16	89.98(3)	In#11-Ce-In#16	107.60(2)
In#12-Ce-In#16	87.78(2)	In#13-Ce-In#16	150.26(4)
In#14-Ce-In#16	52.79(3)	In#15-Ce-In#16	113.23(3)
In#8-Ce-In#16	139.28(2)	In#2-Ce-In#16	49.08(4)
Pd-Ce-In#17	50.15(2)	Pd#9-Ce-In#17	89.98(3)
Pd#10-Ce-In#17	155.46(2)	In#11-Ce-In#17	52.79(3)
In#12-Ce-In#17	150.26(4)	In#13-Ce-In#17	87.78(2)
In#14-Ce-In#17	107.60(2)	In#15-Ce-In#17	113.23(3)
In#8-Ce-In#17	139.28(2)	In#2-Ce-In#17	100.30(5)
In#16-Ce-In#17	80.43(4)	In#18-Pd-In#4	107.29(7)
In#18-Pd-In#17	124.96(3)	In#4-Pd-In#17	74.37(3)
In#18-Pd-In#2	74.37(3)	In#4-Pd-In#2	124.96(3)
In#17-Pd-In#2	150.32(9)	In#18-Pd-In#3	74.37(3)
In#4-Pd-In#3	124.96(3)	In#17-Pd-In#3	63.07(4)
In#2-Pd-In#3	108.77(5)	<b>In#18-Pd-In#1</b> 6	124.96(3)
In#4-Pd-In#16	74.37(3)	In#17-Pd-In#16	108.77(5)
In#2-Pd-In#16	63.07(4)	In#3-Pd-In#16	150.32(9)
In#18-Pd-Ce	126.35(4)	In#4-Pd-Ce	126.35(4)
In#17-Pd-Ce	75.16(4)	In#2-Pd-Ce	75.16(4)
In#3-Pd-Ce	75.16(4)	In#16-Pd-Ce	75.16(4)
In#18-Pd-Ce#19	65.13(3)	In#4-Pd-Ce#19	65.13(3)
In#17-Pd-Ce#19	138.99(5)	In#2-Pd-Ce#19	66.97(3)
In#3-Pd-Ce#19	138.99(5)	In#16-Pd-Ce#19	66.97(3)
Ce-Pd-Ce#19	135.19(3)	In#18-Pd-Ce#20	65.13(3)
In#4-Pd-Ce#20	65.13(3)	In#17-Pd-Ce#20	66.97(3)
In#2-Pd-Ce#20	138.99(5)	In#3-Pd-Ce#20	66.97(3)
In#16-Pd-Ce#20	138.99(5)	Ce-Pd-Ce#20	135.19(3)
Ce#19-Pd-Ce#20	89.62(6)		
Ce#17-FU-Ce#20	03.02(0)		

## Symmetry transformations used to generate equivalent atoms:

```
#1 x,y,z+1 #2 -x-1/2,-y+1/2,-z+1 #3 -x+1/2,-y+1/2,-z+1
#4 x,y,-z+3/2 #5 -x,-y,-z+2 #6 x+1/2,y-1/2,z+1
#7 x-1/2,y-1/2,z+1 #8 -x,-y+1,-z+1 #9 x+1/2,y+1/2,z
#10 x-1/2,y+1/2,z #11 x+1/2,y+1/2,-z+3/2 #12 x-1/2,y+1/2,z-1
#13 x+1/2,y+1/2,z-1 #14 x-1/2,y+1/2,-z+3/2 #15 -x,-y+1,z-1/2
#16 -x-1/2,-y+1/2,z-1/2 #17 -x+1/2,-y+1/2,z-1/2 #18 x,y,z-1
#19 x-1/2,y-1/2,z #20 x+1/2,y-1/2,z
```

Office of Naval Research Chemistry Division, Code 385 Defense Technical Information NAWCWD - China Lake Ctr **(2)** China Lake, CA 93555-6001 Building 5, Cameron Station Alexandria, VA 22314 Dr. Peter Seligman (1) NCCOSC - NRAD Dr. James S. Murday San Diego, CA 92152-5000 (1) Chemistry Division, NRL 6100 Dr. Bernard E. Douda Naval Research Laboratory (1) Crane Division Washington, DC 20375-5660 **NAWC** Dr. John Fischer (1) Crane, Indiana 47522-5000 \* Number of copies required 

800 N. Quincy St.

Arlington, VA 22217-5660

\*\*\*\*\*\*\*end page 6\*\*

(1)\*

Technical Report Distribution List

Physical S&T Division - ONR 331

Dr. John C. Pazik